

# 2024 - 2025

# CARP

Carp farming is an ancient form of fish breeding.  
To this day, carp are still farmed for consumption,  
to populate natural waters, and for commercial fish ponds.



Sinking feed



Designed for RAS



Floating feed



Sustainability score



Semi-floating feed



With astaxanthine



Free from land  
animal protein



Low nitrogen and  
phosphorus emission



High digestiblity



Improved resistance



Omega-3 fatty acids



**Alltech® COPPENS**

DEDICATED TO YOUR PERFORMANCE

# AQUATE™

Innovative premix in all **Alltech Coppens'** feeds.

- + Optimizes growth
- + Supports immune response
- + Optimizes digestive function
- + Contributes to mucous barrier protection
- + Contributes to external barrier protection



## BIO-MOS®

is a mannan-oligosaccharide, which is known to bind and drain opportunistic bacteria. This can result in a significant improvement of the intestinal flora. Additionally, it can improve the structure and length of the microvilli in the gut through which the nutrient intake can increase. **BIO-MOS®** contributes to mucous barrier protection.

## ACTIGEN®

is derived from yeast cell walls and supports the fish's immune response. **Actigen®** furthermore optimizes growth in fish.

## BIOPLEX®

is a crucial element in our new premix. **BIOPLEX®** are organically bound trace elements such as zinc, copper, manganese & iron. With **BIOPLEX®** we can improve the health, growth & performance of the fish.

## AQUATE™

### BIO-MOS®

Improves intestinal function

### ACTIGEN®

Helps maintain the immune system

### BIOPLEX®

Improves performance

### TOTAL REPLACEMENT TECHNOLOGY®

Break with tradition and feed your animals the modern way. Alltech has proven that chelated trace minerals in the form of Bioplex® and Sel-Plex® can be included at significantly lower levels while improving animal performance. This optimizes animal mineral requirements and reduces negative environmental impacts. We call this innovation Alltech's Total Replacement Technology™ (TRT).



**Alltech® COPPENS**

# Carp feed overview

Traditional  
farming  
Semi -  
intensive  
Intensive  
RAS

Granulates

Essence

Advance



Micro pellets

Intensiv

Advance



Pre Growers

Intensiv

Premium Select

Pre Grower-15 EF




Growers

Intensiv


Premium Select

Grower-13 EF



Growers with  
AstaX

Grower Astax



Broodstock

Repro

- Artemia replacer
- High survival
- Supports bone development



## COMPOSITION:

### Analyses (%)

Protein	50	Sizes 0.2-0.3 mm
Fat	10	0.3-0.5 mm
Crude fibre	2,1	0.5-0.8 mm
Ash	7,7	
Total P	0,90	
Astaxanthin (mg/kg)	100	

### Vitamins added

Vitamin A (IE/kg)	22500
-------------------	-------

### Energy (MJ/kg)

Gross Energy	20,1
Digestible Energy	17,8

## FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)


Fish weight (g)	Feed size (mm)	< 10 °C	Larvae fed to satiation								According to fish's appetite & O <sub>2</sub> level	
			10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C		
< 0,2	0,2-0,3											
0,2-0,5	0,3-0,5	According to fish's appetite	2,34	2,83	3,42	4,13	5,00	6,05	7,31	8,84	7,96	
0,5-1,5	0,5-0,8		2,01	2,43	2,94	3,55	4,29	5,19	6,28	7,59	6,83	

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

## ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.



- For semi-intensive farming
- High performance
- High survival
- Medium energy starter diet

### COMPOSITION:

Analyses (%)	0.2-0.3 mm	0.3-0.5 mm	0.5-0.8 mm	1.00 mm	1.5 mm
Protein	56	56	56	54	54
Fat	15	15	15	15	15
Crude fibre	0,3	0,3	0,3	0,4	0,4
Ash	11,3	11,3	11,3	9,6	9,6
Total P	1,77	1,77	1,77	1,66	1,66
<b>Vitamins added</b>					
Vitamin A (IE/kg)	16667	16667	16667	11999	11999

### Energy (MJ/kg)

Gross Energy	21,2	21,2	21,2	21,0	21,0
Digestible Energy	19,2	19,2	19,2	19,3	19,3

### FEEDING TABLE FOR OPTIMAL GROWTH


Fish weight (g)	Feed size (mm)	<10°C	>26°C								
			10°C	12°C	14°C	16°C	18°C	20°C	22°C	24°C	26°C
< 0,2	0,2-0,3							Larvae fed to satiation			
0,2-0,5	0,3-0,5		2,23	2,69	3,26	3,94	4,76	5,76	6,96	8,42	7,58
0,5-1,5	0,5-0,8	According to fish's appetite	1,91	2,31	2,80	3,38	4,09	4,94	5,98	7,23	6,51
1,5-5	0,5-0,8		1,66	2,01	2,43	2,94	3,56	4,30	5,21	6,29	5,67
5,0-8,0	1,0-1,5		1,24	1,50	1,82	2,20	2,65	3,21	3,88	4,69	4,23
8,0-15,0	1,5		1,10	1,33	1,61	1,94	2,35	2,84	3,43	4,15	3,74

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- Semi-intensive farming
- Good performance
- Excellent for roach



### COMPOSITION:

Analyses (%)	Sizes
Protein	40
Fat	10
Crude fibre	1,7
Ash	7,4
Total P	1,20
<hr/>	
Vitamins added	
Vitamin A (IE/kg)	13333
<hr/>	
Energy (MJ/kg)	
Gross Energy	19,2
Digestible Energy	16,5

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)

Fish weight (g)	Feed size (mm)	< 10 °C	> 26 °C								
			10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C
8-15	1,5		1,04	1,26	1,52	1,84	2,23	2,69	3,25	3,94	3,54
25-50	2,0	According to fish's appetite	0,87	1,05	1,27	1,53	1,85	2,24	2,71	3,28	2,95


According to fish's appetite & O<sub>2</sub> level

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.



- High energy diet
- High attractivity
- Good performance
- For semi-intensive farming
- Good for autumn feeding

### COMPOSITION:

#### Analyses (%)

Protein	32	Sizes
Fat	15	2.0 mm
Crude fibre	2,2	
Ash	12,4	
Total P	2,13	

#### Vitamins added

Vitamin A (IE/kg)	10002
-------------------	-------

#### Energy (MJ/kg)

Gross Energy	18,9
Digestible Energy	15,8

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)

Fish weight (g)	Feed size (mm)	> 26 °C									
		< 10 °C	10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C
15-25	2.0		1,06	1,28	1,55	1,87	2,26	2,73	3,31	4,00	3,60
25-50	2.0	According to fish's appetite	0,93	1,13	1,36	1,65	1,99	2,41	2,92	3,53	3,17


According to fish's appetite & O2 level

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- Medium energy diet
- For semi-intensive farming
- Good performance



### COMPOSITION:

#### Analyses (%)

Protein	40	Sizes
Fat	10	2.0 mm
Crude fibre	1,7	
Ash	7,4	
Total P	1,20	

#### Vitamins added

Vitamin A (IE/kg)	13333
-------------------	-------

#### Energy (MJ/kg)

Gross Energy	19,2
Digestible Energy	16,5

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)

Fish weight (g)	Feed size (mm)	< 10 °C	> 26 °C								
			10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C
15-25	2,0		1,04	1,26	1,52	1,84	2,23	2,69	3,25	3,94	3,54
25-50	2,0	According to fish's appetite	0,87	1,05	1,27	1,53	1,85	2,24	2,71	3,28	2,95


According to fish's appetite & O2 level

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- High performance
- Very palatable
- Optical feeding control



### COMPOSITION:

#### Analyses (%)

Protein	50	Sizes
Fat	15	
Crude fibre	0,7	
Ash	6,3	
Total P	1,07	

#### Vitamins added

Vitamin A (IE/kg)	11667
-------------------	-------

#### Energy (MJ/kg)

Gross Energy	21,4
Digestible Energy	19,0

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)


Fish weight (g)	Feed size (mm)	< 10 °C According to fish's appetite	> 26 °C							
			10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C
15-25	2,0		0,93	1,13	1,37	1,65	2,00	2,42	2,92	3,53
25-50	2,0		0,82	1,00	1,20	1,46	1,76	2,13	2,58	3,11

According to  
fish's appetite &  
O<sub>2</sub> level

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- High energy diet
- High attractivity
- Good performance
- For semi-intensive farming
- Good for autumn feeding



### COMPOSITION:

#### Analyses (%)

		Sizes
Protein	32	3.0 mm
Fat	15	4.5 mm
Crude fibre	2,2	6.0 mm
Ash	12,4	8.0 mm
Total P	2,13	14.0 mm 20.0 mm

#### Vitamins added

Vitamin A (IE/kg)	10002
-------------------	-------

#### Energy (MJ/kg)

Gross Energy	18,9
Digestible Energy	15,8

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)

Fish weight (g)	Feed size (mm)	< 10 °C According to fish's appetite	> 26 °C								
			10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C
50-100	3,0		0,83	1,00	1,21	1,47	1,77	2,14	2,59	3,14	2,82
100-250	4,5		0,65	0,78	0,94	1,14	1,38	1,67	2,02	2,44	2,20
250-500	6,0		0,55	0,67	0,81	0,98	1,19	1,43	1,73	2,10	1,89
500-1000	6,0		0,48	0,58	0,71	0,85	1,03	1,25	1,51	1,82	1,64
1000-2000	8,0		0,42	0,51	0,61	0,74	0,90	1,09	1,31	1,59	1,43


According to fish's appetite & O2 level

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- Medium energy diet
- For semi-intensive farming
- Good performance



### COMPOSITION:

#### **Analyses (%)**

		<b>Sizes</b>
Protein	40	3.0 mm
Fat	10	4.5 mm
Crude fibre	2,3	6.0 mm
Ash	10,9	9.0 mm
Total P	1,91	

#### **Vitamins added**

Vitamin A (IE/kg)	10004
-------------------	-------

#### **Energy (MJ/kg)**

Gross Energy	18,6
Digestible Energy	14,8

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)

Fish weight (g)	Feed size (mm)	< 10 °C According to fish's appetite	> 26 °C								
			10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C
50-100	3,0		0,84	1,02	1,23	1,49	1,80	2,18	2,63	3,19	2,87
100-250	4,5		0,66	0,79	0,96	1,16	1,40	1,70	2,05	2,48	2,23
250-500	4,5		0,56	0,68	0,82	1,00	1,20	1,46	1,76	2,13	1,92
500-1000	6,0		0,49	0,59	0,72	0,87	1,05	1,27	1,53	1,85	1,67
1000-2000	9,0		0,43	0,52	0,62	0,75	0,91	1,10	1,33	1,61	1,45


According to fish's appetite & O<sub>2</sub> level

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- Semi-intensive farming
- Good performance
- Very palatable
- Optical feeding control



### COMPOSITION:

#### Analyses (%)

Protein	42
Fat	13
Crude fibre	3,3
Ash	10,0
Total P	1,03

#### Sizes

3.0 mm
4.5 mm
6.0 mm

#### Vitamins added

Vitamin A (IE/kg)	10000
-------------------	-------

#### Energy (MJ/kg)

Gross Energy	19,5
Digestible Energy	15,8

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)


Fish weight (g)	Feed size (mm)	< 10 °C	According to fish's appetite & O2 level								
			10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C
50-100	3,0		0,80	0,97	1,18	1,42	1,72	2,08	2,51	3,04	2,74
100-250	4,5		0,63	0,76	0,91	1,11	1,34	1,62	1,96	2,37	2,13
250-500	6,0		0,54	0,65	0,79	0,95	1,15	1,39	1,68	2,03	1,83
500-1000	6,0		0,47	0,57	0,68	0,83	1,00	1,21	1,46	1,77	1,59
1000-2000	6,0		0,41	0,49	0,60	0,72	0,87	1,05	1,27	1,54	1,39

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- For semi-intensive farming
- Good performance
- Optical feeding control



### COMPOSITION:

#### **Analyses (%)**

Protein	38	4.5 mm
Fat	8	6.0 mm
Crude fibre	2,8	8.0 mm
Ash	12,0	
Total P	1,85	
Astaxanthin (mg/kg)	25	

#### **Vitamins added**

Vitamin A (IE/kg)	10000
-------------------	-------

#### **Energy (MJ/kg)**

Gross Energy	17,7
Digestible Energy	14,4

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)


Fish weight (g)	Feed size (mm)	< 10 °C	10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C	> 26 °C
100-250	4.5		0,69	0,83	1,01	1,22	1,47	1,78	2,16	2,61	2,35	
250-500	6.0	According to fish's appetite	0,59	0,72	0,87	1,05	1,27	1,53	1,85	2,24	2,01	
500-1000	6.0		0,52	0,62	0,75	0,91	1,10	1,33	1,61	1,95	1,75	
1000-2000	8.0		0,45	0,54	0,66	0,79	0,96	1,16	1,40	1,70	1,53	

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- Broodstock diet
- Optimal egg development
- High egg quality and fry survival



## COMPOSITION:

### Analyses (%)

		Sizes
Protein	48	6.0 mm
Fat	15	9.0 mm
Crude fibre	1,3	
Ash	10,5	
Total P	1,61	
Astaxanthin (mg/kg)	40	

### Vitamins added

Vitamin A (IE/kg)	25000
-------------------	-------

### Energy (MJ/kg)

Gross Energy	20,2
Digestible Energy	17,6

## FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)

Fish weight (g)	Feed size (mm)	< 10 °C	> 26 °C								
			10 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	
> 1500	6.0/9.0	According to fish's appetite	0,35	0,43	0,52	0,63	0,76	0,91	1,11	1,34	1,20


According to fish's appetite & O2 level

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

## ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.