

2024 - 2025

SEABASS & SEABREAM

Marine fish farming is characterized by its professionalism and rapid development. We are front runners in this field, developing trendsetting fish feed for European bass and sea bream.



Sinking feed



Designed for RAS



Floating feed



Sustainability score



Semi-floating feed



With astaxanthine



Free from land animal protein



Low nitrogen and phosphorus emission



High digestibility



Improved resistance



Omega-3 fatty acids



SEABASS &
SEABREAM

AQUATE™

Innovative premix in all **Alltech Coppens'** feeds.

- + Optimizes growth
- + Supports immune response
- + Optimizes digestive function
- + Contributes to mucous barrier protection
- + Contributes to external barrier protection



BIO-MOS®

is a mannan-oligosaccharide, which is known to bind and drain opportunistic bacteria. This can result in a significant improvement of the intestinal flora. Additionally, it can improve the structure and length of the microvilli in the gut through which the nutrient intake can increase. **BIO-MOS®** contributes to mucous barrier protection.

BIO-MOS®

Improves intestinal function

BIOPLEX®

is a crucial element in our new premix. **BIOPLEX®** are organically bound trace elements such as zinc, copper, manganese & iron. With **BIOPLEX®** we can improve the health, growth & performance of the fish.

BIOPLEX®

Improves performance

Break with tradition and feed your animals the modern way.

Alltech has proven that chelated trace minerals in the form of Bioplex® and Sel-Plex® can be included at significantly lower levels while improving animal performance. This optimizes animal mineral requirements and reduces negative environmental impacts. We call this innovation Alltech's Total Replacement Technology™ (TRT).

TOTAL REPLACEMENT TECHNOLOGY®

ACTIGEN®

is derived from yeast cell walls and supports the fish's immune response. **Actigen®** furthermore optimizes growth in fish.

ACTIGEN®



Helps maintain the immune system

AQUATE™



Alltech® COPPENS

Marine feed overview



- For semi-intensive farming
- High performance
- High survival
- Medium energy starter diet



COMPOSITION:

Analyses (%)	0.2-0.3 mm	0.3-0.5 mm	0.5-0.8 mm	1.0 mm	1.5 mm
Protein	56	56	56	54	54
Fat	15	15	15	15	15
Crude fibre	0,3	0,3	0,3	0,4	0,4
Ash	11,3	11,3	11,3	9,6	9,6
Total P	1,77	1,77	1,77	1,66	1,66

Vitamins added

Vitamin A (IE/kg)	16667	16667	16667	11999	11999
-------------------	-------	-------	-------	-------	-------

Energy (MJ/kg)

Gross Energy	21,2	21,2	21,2	21,0	21,0
Digestible Energy	19,2	19,2	19,2	19,3	19,3


FEEDING TABLE FOR OPTIMAL GROWTH

Fish weight (g)	Feed size (mm)	< 12 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C	28 °C	> 28 °C
< 0.1	0.2-0.3		Larvae fed to satiation									
0.1-0.2	0.3-0.5	According to fish's appetite	1,53	1,95	2,60	3,45	4,59	6,11	8,51	7,41	5,93	According to fish's appetite & O2 level
0.2-0.4	0.5-0.8		1,41	1,80	2,39	3,18	4,23	5,63	7,85	6,83	5,46	
0.4-1.0	0.5-0.8		1,19	1,52	2,02	2,69	3,57	4,75	6,63	5,77	4,61	
1-4	1.0		0,92	1,18	1,57	2,08	2,77	3,68	5,14	4,47	3,58	
4-8	1.0-1.5		0,77	0,99	1,32	1,75	2,33	3,09	4,31	3,75	3,00	
8-15	1.5		0,68	0,87	1,15	1,54	2,04	2,72	3,79	3,29	2,64	

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- Semi-intensive farming
- Good performance



COMPOSITION:

Analyses (%)		Sizes
Protein	45	2.0 mm
Fat	18	
Crude fibre	1,2	
Ash	9,0	
Total P	1,42	

Vitamins added	
Vitamin A (IE/kg)	11667

Energy (MJ/kg)	
Gross Energy	21,3
Digestible Energy	18,5

FEEDING TABLE FOR OPTIMAL GROWTH


Fish weight (g)	Feed size (mm)	< 12 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C	28 °C	> 28 °C
15-25	2.0	According to fish's appetite	0,46	0,59	0,79	1,05	1,39	1,85	2,59	3,28	2,62	According to fish's appetite & O2 level
25-50	2.0		0,41	0,52	0,70	0,93	1,23	1,64	2,28	2,89	2,31	

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- High protein level
- High performance
- Also very good for perch and pike perch



COMPOSITION:

Analyses (%)		Sizes
Protein	54	2.0 mm
Fat	15	
Crude fibre	0,8	
Ash	9,9	
Total P	1,56	

Vitamins added

Vitamin A (IE/kg)	11667
-------------------	-------

Energy (MJ/kg)

Gross Energy	20,8
Digestible Energy	18,4

FEEDING TABLE FOR OPTIMAL GROWTH


Fish weight (g)	Feed size (mm)	< 12 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C	28 °C	> 28 °C
15-25	2.0		0,47	0,61	0,81	1,07	1,43	1,90	2,65	3,36	2,69	
25-50	2.0	According to fish's appetite	0,42	0,54	0,71	0,95	1,26	1,68	2,34	2,96	2,37	According to fish's appetite & O2 level

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.

- High energy diet
- Fast and efficient growth
- Especially good for bass



COMPOSITION:

Analyses (%)

Protein	43 - 45	Sizes	3.0 mm
Fat	20 - 23		4.5 mm
Crude fibre	1 - 2		6.0 mm
Ash	4 - 8		
Total P	0,98		

Vitamins added

Vitamin A (IE/kg)	9138
-------------------	------

Energy (MJ/kg)

Gross Energy	21,4 - 23,4
Digestible Energy	19,0 - 19,3

FEEDING TABLE FOR OPTIMAL GROWTH


Fish weight (g)	Feed size (mm)	< 12 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C	28 °C	> 28 °C
50-100	3.0		0,60	0,73	0,88	1,06	1,29	1,56	1,88	2,28	1,82	
100-200	3.0	According to fish's appetite	0,48	0,58	0,71	0,85	1,03	1,25	1,51	1,83	1,46	According to fish's appetite & O2 level
200-300	4.5		0,44	0,53	0,64	0,77	0,93	1,13	1,36	1,65	1,32	
300-500	4.5		0,39	0,47	0,57	0,69	0,83	1,00	1,21	1,47	1,17	
> 500	4.5		0,37	0,44	0,54	0,65	0,78	0,95	1,14	1,38	1,11	

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.



- Broodstock diet
- Optimal egg development
- High egg quality and fry survival

COMPOSITION:

Analyses (%)		Sizes
Protein	48	
Fat	15	9.0 mm
Crude fibre	1,3	
Ash	10,5	
Total P	1,61	
Astaxanthin (mg/kg)	40	

Vitamins added

Vitamin A (IE/kg)	25000
-------------------	-------

Energy (MJ/kg)

Gross Energy	20,2
Digestible Energy	17,6

FEEDING TABLE FOR OPTIMAL GROWTH

Fish weight (g)	Feed size (mm)	< 12 °C	12 °C	14 °C	16 °C	18 °C	20 °C	22 °C	24 °C	26 °C	28 °C	> 28 °C
> 600	9,0		0,20	0,24	0,29	0,35	0,43	0,51	0,62	0,75	0,60	

According to fish's appetite


According to fish's appetite & O2 level

* The feeding advice is expressed in % biomass/day.

* This feeding table is a guideline only and based on optimal conditions.

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.