

- Designed for semi-intensive farming
- Good performance
- High flesh quality



### COMPOSITION:

#### Analyses (%)

		Sizes
Protein	41 - 44	4.5 mm
Fat	20 - 23	6.0 mm
Crude fibre	1,5 - 2,5	8.0 mm
Ash	4 - 8	
Total P	0.86	
Astaxanthin (mg/kg)	60.0	

#### Vitamins added

Vitamin A (IE/kg)	9067
-------------------	------

#### Energy (MJ/kg)

Gross Energy	21,4 - 23,4
Digestible Energy	19,2 - 19,5
Net energy	14.4

### FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR):

Fish weight (g)	Feed size (mm)	< 6 °C	6 °C	8 °C	10 °C	12 °C	14 °C	16 °C	18 °C	> 18 °C
100-200	4.5		0.61	0.74	0.89	1.08	1.30	1.57	1.27	
200-300	4.5		0.55	0.66	0.80	0.97	1.17	1.42	1.15	
300-400	4.5	according	0.51	0.62	0.75	0.91	1.10	1.33	1.08	according to fish's appetite and O2 level
400-500	6.0	to fish's	0.49	0.59	0.71	0.86	1.04	1.26	1.02	
500-750	6.0	appetite	0.46	0.56	0.67	0.81	0.99	1.19	0.97	
750-1000	6.0/8.0		0.43	0.52	0.63	0.76	0.92	1.11	0.90	

### FEEDING TABLE FOR OPTIMAL GROWTH


Fish weight (g)	Feed size (mm)	< 6 °C	6 °C	8 °C	10 °C	12 °C	14 °C	16 °C	18 °C	> 18 °C
100-200	4.5		0.79	0.96	1.16	1.40	1.69	2.04	1.66	
200-300	4.5		0.71	0.86	1.04	1.26	1.53	1.85	1.50	
300-400	4.5	according	0.67	0.81	0.98	1.18	1.43	1.73	1.40	according to fish's appetite and O2 level
400-500	6.0	to fish's	0.63	0.77	0.93	1.12	1.36	1.64	1.33	
500-750	6.0	appetite	0.60	0.72	0.88	1.06	1.28	1.55	1.26	
750-1000	6.0/8.0		0.56	0.68	0.82	0.99	1.19	1.44	1.17	

\* The feeding advice is expressed in % biomass/day.

\* This feeding table is a guideline only and based on optimal conditions.

### ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.