

- Designed for RAS and intensive farming
- High performance
- Very suitable for char and hybrids
- Beneficial DP:DE ratio
- High energy phase feeding diet



COMPOSITION:

Analyses (%)	3.0 mm	4.5 mm
Protein	43 - 45	42 - 44
Fat	24 - 27	26 - 29
Crude fibre	4 - 8	4 - 8
Ash	1 - 2	1 - 2
Total P	0,89	0,86

Vitamins added

Vitamin A (IE/kg)	9678	9678
-------------------	------	------

Energy (MJ/kg)

Gross Energy	22,5 - 24,5	23,0 - 25,0
Digestible Energy	20,4 - 20,8	20,9 - 21,3
Net energy	15,7	16,2

FEEDING TABLE FOR LOW FEED CONVERSION RATIO (FCR)


Fish weight (g)	Feed size (mm)	< 6 °C	6 °C	8 °C	10 °C	12 °C	14 °C	16 °C	18 °C	> 18 °C
35-100	3.0	According to fish's appetite	0,72	0,87	1,05	1,27	1,54	1,86	1,51	According to fish's appetite & O2 level
100-200	3.0/4.5		0,58	0,70	0,84	1,02	1,23	1,49	1,21	
200-300	4.5		0,52	0,63	0,76	0,92	1,11	1,35	1,09	
300-400	4.5		0,49	0,59	0,71	0,86	1,04	1,26	1,02	
400-500	4.5		0,46	0,56	0,68	0,82	0,99	1,20	0,97	
500-750	4.5		0,44	0,53	0,64	0,77	0,94	1,13	0,92	
750-1000	4.5		0,41	0,49	0,60	0,72	0,87	1,05	0,85	

FEEDING TABLE FOR OPTIMAL GROWTH

Fish weight (g)	Feed size (mm)	< 6 °C	6 °C	8 °C	10 °C	12 °C	14 °C	16 °C	18 °C	> 18 °C
35-100	3.0	According to fish's appetite	1,04	1,26	1,52	1,84	2,23	2,70	2,18	According to fish's appetite & O2 level
100-200	3.0/4.5		0,84	1,01	1,22	1,48	1,79	2,16	1,75	
200-300	4.5		0,76	0,91	1,10	1,34	1,62	1,95	1,58	
300-400	4.5		0,71	0,85	1,03	1,25	1,51	1,83	1,48	
400-500	4.5		0,67	0,81	0,98	1,19	1,44	1,74	1,41	
500-750	4.5		0,63	0,77	0,93	1,12	1,36	1,64	1,33	
750-1000	4.5		0,59	0,72	0,87	1,05	1,26	1,53	1,24	

ECOLOGICAL FIGURES:

Discharge per 1000 kg production



The values of the nutrients and vitamins are from the time of writing.

These values can vary due to natural variation in the ingredients. We reserve the right to change our recipe.

For the exact values we refer to the label.